

← Patientenetikette hier ▲ positionieren!

Name	Geschlecht	
Vorname	Geburtsdatum	
Straße / Nr.		
PLZ / Ort		
Allg. Geb.-Kl. <input type="checkbox"/>	Sonder Geb.-Kl. <input type="checkbox"/>	Selbstzahler <input type="checkbox"/>
Klin. Hinweis bzw. Diagnose		

Information for admission as an inpatient

Dear patient,

Your admission as an inpatient has been planned for _____ at _____ .

- You may eat as usual on the day of admission, i.e. an empty stomach is not necessary at the time of your admission.
- Nil by mouth from midnight onwards, i.e. no food, drink, smoking or chewing gum.

Important: Ensure you remove piercings, jewellery, make-up and nail polish before your operation.

Please register as an inpatient in the 'Frauen- und Kopfklinik' building no. 3

- at the central office of the _____ ward on the ____ floor
- at the Gynaecology Day Unit 4 (Tagesklinik Gynäkologie 4) on the 6th floor 'Nord'
NB: Ensure you arrange for someone to pick you up.
- at the Oncological Clinic (Onkologische Sprechstunde) on the 2nd floor

As an inpatient please ensure you take the following items with you:

- Any existing medical results/findings (internist, neurologist, anaesthesia outpatient clinic) and documentation (consent to surgery)
- Bathrobe, slippers, pyjamas, towels and toiletries
- A sufficient quantity of any medication that you take on a regular basis

If you are unable to keep the arranged admission appointment, please contact us as early as possible by telephone to cancel or change your admission:

- General Outpatient Clinic050 504-230 60 (Mon – Fri 8.00 am – 12.00 noon)
- Consultant's Surgery050 504-230 56 (Mon – Thurs 10.00 am – 4.00 pm)
- Oncological Clinic050 504-230 64

The team of nursing staff and doctors at
Innsbruck University Hospital's Department of Gynaecology and Obstetrics
wish you a comfortable stay and a speedy recovery.

Important!

During the 7 days before your surgery it is important that you **do not take any blood thinning medication** (e.g. Sintrom, Thrombo-Ass, Aspirin or any medication containing acetylsalicylic acid).

Talk to your doctor in the outpatient department about switching to alternative medication. The intake of oral antidiabetic drugs or the administration of insulin must be agreed upon with the outpatient physician.